

Post Test --- Insulin pumps and the Diabetic Patient

1. Insulin pumps can only be worn in specific locations?
 - a. True
 - b. False

2. The patient must go to a pharmacy, hospital, or physicians office to refill their insulin pump?
 - a. True
 - b. False

3. How often should a patient with an insulin pump check their blood sugars?
 - a. 3 times a day
 - b. 3 times a week
 - c. Twice a day in the morning and at night
 - d. At least 4 times a day

4. Which of the following is not a component of the insulin pump?
 - a. A pump reservoir filled with insulin
 - b. A tube going from the pump to the patient
 - c. A battery operated pump
 - d. A computer chip that controls the amount of insulin administered

5. The insulin pump delivers a set amount of insulin continuously throughout the day, this is called the _____ rate.
 - a. Set
 - b. Continuous
 - c. Bolus
 - d. Basal

6. The patient can set the pump to deliver a preset amount of insulin at a set time during the day. This is called the _____ rate.
 - a. Set
 - b. Preset
 - c. Bolus
 - d. Basal

7. The only approved site for infusion of insulin to the patient is in the abdomen?
 - a. True
 - b. False

8. Which method do the manufacturers least recommend for resolving pump issues?
 - a. Suspend the pump
 - b. Remove the batteries
 - c. Disconnect the tubing
 - d. completely remove to pump from the patient